

## **CURRENT ASPECTS OF EPIDEMIOLOGICAL AND CLINICAL ISSUES OF EPSTEIN-BARR VIRUS INFECTION IN CHILDREN**

***Rabbimova Nodira Tashtemirovna***

*PhD, Assistant of Department of Infectious Diseases,*

*Samarkand State Medical University,*

*Samarkand, Uzbekistan*

**ABSTRACT.** The article presents information on the clinical course of infectious mononucleosis in 142 patients aged 1 to 14 years living in the Samarkand region. It was found that at the current stage, the disease retains all the features of its clinical course. If the diagnosis of infectious mononucleosis is made in time and accurately, the effectiveness of treatment in infected patients is shown by reducing the duration of fever, the severity of the proliferative syndrome in the lymph nodes, and the faster cessation of the cytolysis process.

**Keywords:** atypical mononuclear cells, children, Epstein-Barr virus (EBV), infectious mononucleosis.

### **АКТУАЛЬНЫЕ АСПЕКТЫ ЭПИДЕМИОЛОГИЧЕСКИХ И КЛИНИЧЕСКИХ ВОПРОСОВ ЭПШТЕЙН-БАРР ВИРУСНОЙ ИНФЕКЦИИ У ДЕТЕЙ**

**АБСТРАКТ.** В статье представлены сведения о клиническом течении инфекционного мононуклеоза у 142 больных в возрасте от 1 до 14 лет, проживающих в Самаркандской области. Установлено, что на современном этапе заболевание сохраняет все особенности клинического течения. При своевременной и точной постановке диагноза инфекционного мононуклеоза эффективность лечения инфицированных больных проявляется уменьшением продолжительности лихорадки, выраженности пролиферативного синдрома в лимфатических узлах, более быстрым прекращением процесса цитолиза.

**Ключевые слова:** атипичные мононуклеары, дети, вирус Эпштейна-Барр (ВЭБ), инфекционный мононуклеоз.

## **BOLALARDA EPSHTEYN-BARR VIRUSLI KASALLIGINING KLINIK VA EPIDEMIOLOGIK KECHISH ASPEKTLARI**

**ABSTRACT.** Maqolada Samarqand viloyatida yashovchi 1 yoshdan 14 yoshgacha bo'lgan 142 nafar bemorda yuqumli mononuklyozning klinik kechishi haqida ma'lumotlar keltirilgan. Hozirgi bosqichda kasallik o'zining klinik kechishining barcha xususiyatlarini saqlab qolganligi aniqlandi. Yuqumli mononuklyozni o'z vaqtida va to'g'ri tashxislash bilan kasallangan bemorlarni davolash samaradorligi isitma davomiyligining pasayishi, limfa tugunlarida proliferativ sindromning og'irligi va sitoliz jarayonining tezroq to'xtashi bilan namoyon bo'ladi.

**Kalit so'zlar:** atipik mononuklear hujayralar, bolalar, Epstein-Barr virusi (EBV), yuqumli mononuklyoz.

**Relevance.** Epstein-Barr virus infectious mononucleosis is one of the main infectious diseases in childhood [3,5,7,11]. In most children, the disease ends with a good outcome, without complications, but the Epstein-Barr virus can remain in the body of an infected person for life. In a small number of children, the preservation of the virus in such a state can serve as a stimulus for the development of processes such as lymphoproliferative syndrome, oncological pathological changes, chronic fatigue syndrome, and hemophagocytic syndromes [7,9,12].

Currently, the known Epstein-Barr virus (herpes virus type IV) does not always serve as an etiological factor of acute infectious mononucleosis; however, it accounts for 90-95% of all cases. In the remaining cases, the disease may be caused by other types of viruses such as human immunodeficiency virus, adenovirus, cytomegalovirus, and herpes viruses of type VI [2,10,13]. According to the ICD, infectious mononucleosis is classified depending on the causative agents of the disease as follows: Epstein-Barr virus (B27.0); cytomegalovirus (CMV) (B27.1); infectious mononucleosis of other etiology (B27.8); infectious mononucleosis of unknown etiology (B27.9) if the etiological cause of the disease is not determined in patients with clinical symptoms characteristic of infectious mononucleosis.

Etiotropic treatment for infectious mononucleosis has not been developed to date. Treatment of infectious mononucleosis in the acute period aims to reduce the replication of the Epstein-Barr virus and to induce an adequate immune response [4, 6,8,14,15].

**The aim** of this study is to investigate the specific clinical features of infectious mononucleosis of Epstein-Barr virus etiology in children residing in the Samarkand region.

**Methods and Materials:** Comparative and clinical examinations, general and biochemical analysis of biomaterials, and PCR examination methods were conducted on patients treated at the Samarkand Regional Infectious Diseases Clinical Hospital from 2007 to 2023. A total of 142 patients aged 1 to 14 years diagnosed with infectious mononucleosis were monitored.

**Results and Discussion:** Analysis of the obtained results revealed that infectious mononucleosis exhibits clear seasonality. Throughout the year, the incidence of this disease was primarily recorded in the winter and spring months (24% and 46.0%, respectively), with a lower percentage of cases occurring in other seasons (30%). (Fig. 1).




Figure 1. Seasonality of infectious mononucleosis

When the gender incidence of the disease was studied, the ratio of boys to girls was analyzed. The percentage of boys affected was 51%, while the percentage of girls affected was 49%( Figure 2).





Figure 2. Gender distribution of the disease

In the course of our research, we discovered that 28.9% of patients diagnosed with infectious mononucleosis arrived at the hospital late. This delay may be attributed to the clinical symptoms of infectious mononucleosis being masked by the symptoms of coexisting diseases.

When evaluating the clinical course of the disease, the severity was categorized as follows: mild course - 18.2%, moderate to severe course - 76.7%, severe course - 5.1% (Figure 3).



Picture 3. Disease severity degrees according to distribution

In the clinical course of the disease, the main clinical signs observed in the past include the following pointers: rhinitis (96%), elevation of body temperature from subfebrile to febrile degree (91.4%), enlargement of the liver and spleen (91.7%),

tonsillitis (monocytic angina) (84.2%), lymph node hyperplasia (67.2%), adenoiditis (11.5%), and exanthema (18.7%) (Fig. 4).




Figure 4. Clinical signs of the disease.

In the analytical discussion of the diagnostic basis of the disease, general blood, urine, and stool analyses, as well as blood biochemical analysis, were conducted. Lymphocytic leukocytosis and atypical mononuclear cells were identified from hematological changes, and PCR testing was performed to detect Epstein-Barr virus DNA in the blood for verification. Chest X-rays were carried out using instrumental examination methods as per protocol, and all patients underwent ultrasound examination.

All patients received treatment according to standard care protocols, including detoxification therapy, symptomatic relief, and antibiotic therapy guided by clinical instructions. Recombinant interferon inducers were administered to stimulate the immune system. Dosages of all medications were adjusted based on the patient's age, body weight, and recommended dosages in the instructions.

**Conclusion:** Our research revealed that the seasonality of the disease primarily corresponds to the winter and spring months. Infectious mononucleosis retains all its clinical features at the present stage. Fever, runny nose, and hepatosplenomegaly were identified as prominent clinical signs. Early diagnosis of the disease plays a crucial role in reducing symptoms of cytolytic syndrome and mesenchymal inflammation.

### References:

1. Infectious mononucleosis /MV Krasnov, IA Stekolshikova, MG Borovkova, LV Andreeva //Sovremennye problemi nauki i obrazovaniya.-2015.-№ 2.- S. 24-26.
2. Infectious mononucleosis and detection, associated with herpes virus 4 and 5 /EB Kasimova, OA Bashkina, XM Galimzyanov, SJ Netalieva //Infectious diseases.-2012-Vol.10, No.3-S.44-47.-Text : neposredstvenniy.
3. Kramar, LV Herpeticheskaya infection and mononucleozopodobniy syndrome and detection /LV Kramar, OA Karpukhina.-Volgograd: Izd-vo VolgGMU, 2016.-292 p.-Text: neposredstvenniy.
4. Kramar, LV Complex therapy of Epstein-Barr virus infection and detey-Text: nepostredstvenniy/LVKramar, OAKarpukhina//Archive of internal medicine. -2012.-No. 1 (3). - S. 25-29.
5. Tuychiev L. N. et al. NASOPHARYNGEAL EXTRACTION OF S. PNEUMONIAE FROM ADULT PATIENTS WITH ACUTE RESPIRATORY INFECTIONS AND ANTIBIOTIC RESISTANCE OF ISOLATED STRAINS //Art of Medicine. International Medical Scientific Journal. – 2022. – Т. 2. – №. 1.
6. Раббимова Н. Т., Матякубова Ф. Э., Тиркашев О. С. ЧАСТОТА ВЫДЕЛЕНИЯ STREPTOCOCCUS PNEUMONIAE ПРИ ОСТРЫХ РЕСПИРАТОРНЫХ ИНФЕКЦИЯХ ДЫХАТЕЛЬНЫХ ПУТЕЙ //VOLGAMEDSCIENCE. – 2021. – С. 589-591.
7. Tuychiev L. N. et al. Antimicrobial susceptibility OF S. Pneumoniae, isolated from adults //湖南大学学报 (自然科学版). – 2021. – Т. 48. – №. 11.
8. Раббимова Н. и др. Математическое моделирование и прогнозирование заболеваемости кожным лейшманиозом в республике узбекистан //Журнал проблемы биологии и медицины. – 2017. – №. 1 (93). – С. 104-107.
9. Сувонкулов У. и др. Идентификация видовой принадлежности возбудителей кожного лейшманиоза методом полимеразной цепной реакции //Журнал проблемы биологии и медицины. – 2016. – №. 3 (89). – С. 91-92.
10. Egamovna M. F. et al. CLINICAL AND EPIDEMIOLOGICAL FEATURES OF THE COURSE OF SHIGELLOSIS IN ADULTS AT THE PRESENT STAGE IN 2009-2019 //Web of Scientist: International Scientific Research Journal. – 2022. – Т. 3. – №. 5. – С. 1285-1294.
11. Абдухалилова Г. К. и др. Назофарингеальное носительство str. e у взрослых. – 2022.

Egamovna M. F. et al. CLINICAL AND EPIDEMIOLOGICAL FEATURES OF THE COURSE OF SHIGELLOSIS IN ADULTS AT THE PRESENT STAGE IN 2009-2019 //Web of Scientist: International Scientific Research Journal. – 2022. – Т. 3. – №. 5. – С. 1285-1294.

12. Абдухалилова Г. К. и др. Динамика устойчивости к антибиотикам и частота назофарингеального выделения *S. Pneumoniae* у взрослых с острыми респираторными инфекциями. – 2022.

13. Ярмухамедова Н. и др. Особенности течения хронического гепатита с на фоне туберкулеза //Журнал вестник врача. – 2019. – Т. 1. – №. 1. – С. 129-132.1

14. Anvarovna, Y. N., Egamovna, M. F., Tashtemirovna, R. N., Buribayevna, M. G., & Saidovich, T. O. (2021). Clinical and Epidemiological Characteristics of Shigellosis in Adults at the Contemporary Stage. *Central Asian Journal of Medical and Natural Science*, 2(3), 311-318. <https://doi.org/10.47494/cajmns.v2i3.221>

15. Тиркашев, О. С. Клинико-эпидемиологическая характеристика кори в Самаркандской области / О. С. Тиркашев, Ф. Э. Матякубова, Н. Т. Раббимова // VOLGAMEDSCIENCE : Сборник тезисов VII Всероссийской конференции молодых ученых и студентов с международным участием: материалы конференции, Нижний Новгород, 16–18 марта 2021 года. – Нижний Новгород: Федеральное государственное бюджетное образовательное учреждение высшего образования "Приволжский исследовательский медицинский университет" Министерства здравоохранения Российской Федерации, 2021. – С. 624-625. – EDN GZYHJQ.

16. Tirkashev O. S. et al. MEASLES AT THE PRESENT STAGE //Web of Scientist: International Scientific Research Journal. – 2022. – Т. 3. – №. 5. – С. 177-185.